



Rutter, G. A., McCormack, J. G., Halestrap, A. P., & Denton, R. M. (2020). The roles of cytosolic and intramitochondrial Ca^{2+} and the mitochondrial Ca^{2+} -uniporter (MCU) in the stimulation of mammalian oxidative phosphorylation. *Journal of Biological Chemistry*, (2020) 295(30), 10506-10506. <https://doi.org/10.1074/jbc.L120.013975>

Peer reviewed version

Link to published version (if available):
[10.1074/jbc.L120.013975](https://doi.org/10.1074/jbc.L120.013975)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via American Society for Biochemistry and Molecular Biology at <https://www.jbc.org/content/295/30/10506.extract> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

The roles of cytosolic and intramitochondrial Ca^{2+} and the mitochondrial Ca^{2+} -uniporter (MCU) in the stimulation of mammalian oxidative phosphorylation

Guy A. Rutter^{1,2*}, James G. McCormack³, Andrew P. Halestrap⁴ &
Richard M. Denton⁴

¹Section of Cell Biology and Functional Genomics, Division of Diabetes, Endocrinology and Metabolism, Department of Metabolism, Reproduction and Digestion, Imperial College London, London, U.K. and ²Lee Kong Chian School of Medicine, Nan Yang Technological University, Singapore, ³McC+R&D Consulting Ltd., 12 Carlton Terrace, Edinburgh EH7 5DD, UK, ⁴School of Biochemistry, University of Bristol, Bristol BS8 1TD, U.K.

*Correspondence to Professor Guy Rutter, g.rutter@imperial.ac.uk

Szibor et al. (1) conclude that cytosolic $[\text{Ca}^{2+}]$ ($[\text{Ca}^{2+}]_{\text{cyt}}$) activation of the malate-aspartate shuttle is more important than mitochondrial $[\text{Ca}^{2+}]$ ($[\text{Ca}^{2+}]_{\text{mit}}$) activation of intramitochondrial dehydrogenases in the regulation of pyruvate oxidation by mammalian mitochondria. However, the authors employed unphysiological conditions with isolated mitochondria (saturating ADP), with synaptosomes, thymocytes and fibroblasts (uncoupler and high pyruvate) and with perfused hearts (high pyruvate) that would inevitably deliver the results obtained. Pyruvate dehydrogenase (PDH) activity is largely determined by the ratio of active non-phosphorylated PDH to inactive phosphorylated PDH (PDHP); this ratio depends on the activities of the ATP-linked PDH kinase

(inhibited by ADP and pyruvate), and PDHP phosphatase (activated by $[Ca^{2+}]_{mit}$) (2).

Under all the conditions used by Szibor et al. (1), we would predict PDH/PDHP to be so high that further activation by increased $[Ca^{2+}]_{mit}$ is very limited. No measurements of PDH/PDHP ratios were reported. Under physiologically-appropriate conditions, where PDH/PDHP ratios are lower, there is extensive evidence that $[Ca^{2+}]_{mit}$ is an important determinant of pyruvate oxidation (3-5).

Regulation of the malate-aspartate shuttle by $[Ca^{2+}]_{cyt}$ (allowing increased mitochondrial oxidation of cytoplasmic NADH) complements the regulation of intramitochondrial dehydrogenases by $[Ca^{2+}]_{mit}$ (2). We regard the latter as an evolutionary refinement of “intrinsic” mechanisms (also present in lower organisms) enabling increased ATP supply without lowering ATP/ADP ratios (2). Indeed, the MCU blocker ruthenium red decreases ATP/ADP ratios in stimulated hearts (6). The intrinsic mechanisms may operate in MCU-null mice, but these mice do show exercise abnormalities (7). Furthermore, mitochondrial Ca^{2+} influx is not completely suppressed by MCU deletion (7).

254 words including reference numbers

Reference List

1. Szibor, M., Gizatullina, Z., Gainutdinov, T., Endres, T., Debska-Vielhaber, G., Kunz, M., Karavasili, N., Hallmann, K., Schreiber, F., Bamberger, A., Schwarzer, M., Doenst, T., Heinze, H. J., Lessmann, V., Vielhaber, S., Kunz, W. S., and Gellerich, F. N. (2020) Cytosolic, but not matrix, calcium is essential for adjustment of mitochondrial pyruvate supply. *J. Biol. Chem.* **295**, 4383-4397
2. McCormack, J. G., Halestrap, A. P., and Denton, R. M. (1990) Role of calcium ions in regulation of mammalian intramitochondrial metabolism. *Physiol Rev* **70**, 391-425
3. McCormack, J. G., Edgell, N. J., and Denton, R. M. (1982) Studies on the interactions of Ca^{2+} and pyruvate in the regulation of rat heart pyruvate dehydrogenase activity. Effects of starvation and diabetes. *Biochem J* **202**, 419-427
4. Hansford, R. G. and Castro, F. (1985) Role of Ca^{2+} in pyruvate dehydrogenase interconversion in brain mitochondria and synaptosomes. *Biochem. J.* **227**, 129-136

5. Griffiths, E. J. and Rutter, G. A. (2009) Mitochondrial calcium as a key regulator of mitochondrial ATP production in mammalian cells. *Biochim. Biophys. Acta.* **1787**, 1324-1333
6. Unitt, J. F., McCormack, J. G., Reid, D., MacLachlan, L. K., and England, P. J. (1989) Direct evidence for a role of intramitochondrial Ca^{2+} in the regulation of oxidative phosphorylation in the stimulated rat heart Studies using ^{31}P n.m.r. and ruthenium red. *Biochem. J.* **262**, 293-301
7. Pan, X., Liu, J., Nguyen, T., Liu, C., Sun, J., Teng, Y., Fergusson, M. M., Rovira, I. I., Allen, M., Springer, D. A., Aponte, A. M., Gucsek, M., Balaban, R. S., Murphy, E., and Finkel, T. (2013) The physiological role of mitochondrial calcium revealed by mice lacking the mitochondrial calcium uniporter. *Nat. Cell Biol.* **15**, 1464-1472